



Guidelines for CUT FLOWERS for EXPORT

Flower Protocol

Initial Planting from Seed: Apply dry powdered *Seed Coat* at 5 dry ounces per hectare for ample coverage as though dusting the seed.

At Transplanting: Apply *PercPlus* at 2 ounces per gallon (4 liters) of water into the transplant water upon initial setting out of the plants.

After True Leaf Formation: After third (3) true leaf emerges or 4-5 days after transplant: Apply *PercPlus* at 32 Oz/acre (2.4 L/ha) either as a foliar or in drip irrigation. REPEAT full rate within 10-14 days.

Pre-Bloom (Initial Budding): Apply *CropKarb* at 32 Oz/acre (2.4 L/ha). REPEAT in 10 to 14 days. This application may be made either as a foliar or as a Drip irrigation application.

PLEASE NOTE: These Guidelines are what we feel is the best approach to being able to help improve yield and quality of these horticultural crops. Therefore, please know that we, at DeltAg, have no issues should you review this protocol and have questions or concerns of any kind. While there are other DeltAg products that could be used on these crops, our focus is on the primary products that we would expect to generate the most economical responses in most situations.

NOTE: *CropKarb* may be replaced with a tank-mix of *Potassium Plus* @ 32 Oz, *PercPlus* @ 8 to 16 Oz, and *Boron Plus* @ 4 Oz. *Plant Power* may be substituted for *PercPlus* and vice-versa in any application and at the same rate.

Mixing: Always dilute *Potassium Plus* in water first and follow with other ingredients in listed order to avoid excessive foaming.

These **DeltAg Guidelines** have been utilized to improve overall crop health and are not intended to be utilized as a replacement for basic and customary fertilizer applications. **“Other Crop Specific Guidelines are available upon request.”**