



Guidelines for HEMP - GROWN for SEED

Preplant Soil Application: Apply *Soil Solution* @ 24 Oz/acre. This application should be made in the Fall and incorporated or up to planting in the Spring. For No-till or over a growing crop, increase rate to 32 Oz/acre.

Seed Treatment: DRY Hopper Box Seed Treatment: Apply *Seed Coat* @ 2 to 3 Oz (dry weight) per acre. The standard recommended rate for *Seed Coat* is 4 Oz per 50 pounds of seed.

Soil Applied: For pH that is lower than desired: Broadcast apply *Calcium Plus* @ 64 Oz/acre minimum at or just prior to planting and slightly incorporate no deeper than 1 ½ to 2 inches deep.

NOTE: The design of this treatment is to affect the germination zone only. This is an effort to improve early growth and root development. This treatment is considered temporary and is not intended and **WILL NOT** correct soil pH or replace the need for Lime.

Seedling Foliar: Apply *PercPlus* @ 24 Oz/acre broadcast at 2-3 leaves.

Vegetative Foliar: Apply *PercPlus* @ 24 Oz/acre broadcast 7-10 days after seedling above.

Fruiting Foliar: Apply *CropKarb* @ 32 Oz/acre broadcast by the 3rd or 4th week of bloom.

Sizing Foliar: Apply *CropKarb* @ 32 Oz/acre broadcast 10-14 days after Fruiting Foliar above.

NOTE: *Plant Power* may be substituted for *PercPlus* and vice-versa in any application and at the same rate.

Mixing:

These **DeltAg Guidelines** have been utilized to improve overall crop performance and are not intended to be utilized as a replacement for basic and customary fertilizer applications. “**Other Crop Specific Guidelines are available upon request.**”