



Guidelines for TURF

Dormancy Application: Apply *Soil Solution* @ 32 Oz/acre broadcast. This application can be made in the late fall, winter or early spring, before turf starts to green up. **Note:** It may also be applied on green turf but will not be as beneficial for the specific season of application.

Initial Planting DRY Seed Treatment: Apply *Seed Coat* @ 2 Oz (dry weight) per planted acre of seed.

Spring Application: *PercPlus* @ 24 Oz/acre broadcast. May be applied shortly after turf green up or two to three weeks after green up.

Late Spring Application: *CropKarb* @ 32 Oz/acre and *Iron Plus* @ 8 Oz/acre broadcast
Substitution: *PercPlus* @ 16 Oz/acre and *Potassium Plus* @ 32 Oz/acre with the *Iron Plus* @ 8 Oz/acre.

Mid-Summer Application: *CropKarb* @ 32 Oz/acre and *Iron Plus* @ 8 Oz/acre broadcast.
Substitution: *PercPlus* @ 8 Oz/acre and *Potassium Plus* @ 32 Oz/acre with the *Iron Plus* @ 8 Oz/acre.

Late Summer/Early Fall Application: *CropKarb* @ 32 Oz/acre and *Iron Plus* @ 8 Oz/acre broadcast.
Substitution: *PercPlus* @ 24 Oz/acre and *Potassium Plus* @ 16 Oz/acre with the *Iron Plus* @ 8 Oz/acre.

NOTE: *CropKarb* may be replaced with a tank-mix of *Potassium Plus* @ 32 Oz, *PercPlus* @ 8 to 16 Oz, and *Boron Plus* @ 4 Oz. *Plant Power* may be substituted for *PercPlus* and vice-versa in any application and at the same rate.

Mixing: Always dilute *Potassium Plus* in water first and follow with other ingredients in listed order to avoid excessive foaming.

These **DeltAg Guidelines** have been utilized to improve overall crop health and are not intended to be utilized as a replacement for basic and customary fertilizer applications. **“Other Crop Specific Guidelines are available upon request.”**